This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

GIFT OF
the estate of
Prof. Wm. F. Meyer

ASTRONOMY LIBRARY

Digitized by Google

UNIVERSITY OF CALIFORNIA

Uranography

Young

Digitized by Google

Digitized by Google

Digitized by Google

URANOGRAPHY

Digitized by Google

Digitized by Google

URANOGRAPHY

A BRIEF DESCRIPTION OF

THE CONSTELLATIONS VISIBLE IN THE UNITED STATES

WITH

STAR-MAPS, AND LISTS OF OBJECTS OBSERVABLE WITH A SMALL TELESCOPE

BY

C. A. YOUNG, PH.D., LL.D.

LATE PROFESSOR OF ASTRONOMY IN PRINCETON UNIVERSITY

A SUPPLEMENT TO THE AUTHOR'S "ELEMENTS OF ASTRONOMY FOR HIGH SCHOOLS AND ACADEMIES"

GINN AND COMPANY

BOSTON · NEW YORK · CHICAGO · LONDON

ATLANTA · DALLAS · COLUMBUS · SAN FRANCISCO

Digitized by Google

Astronomy
Calcul. to. Clio

ENTERED AT STATIONERS' HALL

COPYRIGHT, 1889, 1897, BY
CHARLES A. YOUNG

COPYRIGHT, 1917, BY
FREDERICK A. YOUNG

ALL RIGHTS RESERVED

PRINTED IN THE UNITED STATES OF AMERICA

331.8

The Athenæum Press
GINN AND COMPANY · PROPRIETORS · BOSTON · U.S.A.

Digitized by Google

QB802
Y6
1917
ASTRONOMY
LIBRARY

PREFACE.

THIS brief description of the constellations was prepared, at the suggestion of a number of teachers, as an integral part of the author's "Elements of Astronomy." It has been thought best, however, for various reasons, to put it into such a form that it can be issued separately, and used if desired in connection with the larger "General Astronomy," or with any other text-book. Since the Uranography also has to be used more or less in the open air at night, many will probably prefer to have it by itself, so that its use need not involve such an exposure of the rest of the text-book. All references marked Astr. are to the articles of the "Elements of Astronomy."

M577279

Digitized by Google

Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY

ALPHABETICAL LIST OF THE CONSTELLATIONS DESCRIBED OR MENTIONED IN THE URANOGRAPHY.

	Article
Andróměda	16
Anser, *see* Vulpécŭla	54
Antínoüs, *see* Aquila	56
Antlia	45
Aquarius	63
Aquila	56
Argo Navis	34
Aries	19
Aurīga	22
Boötes	42
Camelopárdalis	14
Cancer	35
Canes Venatici	41
Canis Major	33
Canis Minor	31
Capricornus	58
Cassiŏpéia	9, 10
Centaurus	45
Cepheus	11
Cetus	20
Coma Berĕnīces	40
Columba	28
Corōna Borealis	43
Corvus	38
Crater	38
Cygnus	53
Delphīnus	59
Draco	12, 13
Equúleus	60
Erídănus	26
Gemini	30
Grus	64
Hércŭles	50, 51
Hydra	38
Lacerta	61
Leo	36
Leo Minor	37
Lepus	27
Libra	44
Lupus	45
Lynx	29
Lyra	52
Monóceros	32
Norma	47
Ophiŭchus	48, 49
Orīon	24, 25
Pégăsus	62
Perseus	21
Phœnix	20
Pisces	17
Piscis Australis	64
(Pleiades)	23
Sagitta	55
Sagittarius	57
Scorpio	46, 47
Sculptor	20
Serpens	48, 49
Serpentarius, *see* Ophiuchus	48, 49
Sextans	37
Taurus	23
Taurus Poniatovii	48
Triángŭlum	18
Ursa Major	5–7
Ursa Minor	8
Virgo	39
Vulpécŭla	54

Digitized by Google

THE GREEK ALPHABET.

Letters.	Name.	Letters.	Name.	Letters.	Name.
A, α,	Alpha.	I, ι,	Iota.	P, ρ, ϱ,	Rho.
B, β,	Beta.	K, κ,	Kappa.	Σ, σ, ς,	Sigma.
Γ, γ,	Gamma.	Λ, λ,	Lambda.	T, τ,	Tau.
Δ, δ,	Delta.	M, μ,	Mu.	Y, υ,	Upsilon.
E, ϵ,	Epsilon.	N, ν,	Nu.	Φ, ϕ,	Phi.
Z, ζ,	Zeta.	Ξ, ξ,	Xi.	X, χ,	Chi.
H, η,	Eta.	O, o,	Omicron.	Ψ, ψ,	Psi.
Θ, θ, ϑ,	Theta.	Π, π,	Pi.	Ω, ω,	Omega.

Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY

URANOGRAPHY,

OR

A Description of the Constellations.

1. A general knowledge of the constellations sufficient to enable one to recognize readily the more conspicuous stars and their principal configurations, is a very desirable accomplishment, and not difficult to attain. It requires of course the actual study of the sky for a number of evenings in different parts of the year; and the study of the sky itself must be supplemented by continual reference to a celestial globe or star-map, in order to identify the stars observed and fix their designations. A well-made globe of sufficient size is the best possible help, because it represents things wholly without distortion, and is easily "rectified" (Astr. 528[1]) for any given hour, so that the stars will all be found in the proper quarter of the (artificial) heavens, and in their true relations. But a globe is clumsy, inconvenient out of doors, and liable to damage; and a good star-map properly used will be found but little inferior in efficiency, and much more manageable.

2. Star-Maps. — Such maps are made on various systems, each presenting its own advantages. None are without more or less distortion, especially near the margin, though they

[1] The references are to the articles in the Author's "Elements of Astronomy," to which this Uranography is a supplement.

Digitized by Google

differ greatly in this respect. In all of them the heavens are represented *as seen from the inside*, and not as on the globe, which represents the sky as if seen from the *outside; i.e.*, the top of the map is *north*, and the east is at the *left* hand; so that if the observer faces the south and holds up the map before and above him, the constellations which are near the meridian will be pretty truly represented.

3. We give a series of four small maps which, though hardly on a large enough scale to answer as a satisfactory celestial atlas, are quite sufficient to enable the student to trace out the constellations and identify the principal stars.

In the map of the *north* circumpolar regions (Map I.), the pole is in the centre, and at the circumference the right-ascension hours are numbered in the same direction as the figures upon a watch face; but with 24 hours instead of 12. The parallels of declination are represented by equidistant and concentric circles. On the three other rectangular maps, which show the equatorial belt of the heavens lying between $+50°$ and $-50°$ of declination, the parallels of declination are equidistant horizontal lines, while the hour-circles are vertical lines also equidistant, but spaced at a distance which is correct for declination 35°, and not at the equator. This keeps the distortion within reasonable bounds even near the margin of the map, and makes it very easy to lay off the place of any object for which the right ascension and declination are given.

The hours of right ascension are indicated on the central horizontal line, which is the equator, and at the top of the map are given the names of the months. *The word* ***September****, for instance, means that the stars which are directly under it upon the map will be near the meridian* ***about nine o'clock*** *in the evening during that month.*

4. The maps show all the stars down to the $4\frac{1}{2}$ magnitude — all that are easily visible on a moonlight night. A few smaller stars are also inserted, where they mark some peculiar configuration or point out some interesting telescopic object. So far as practicable, *i.e.*, north of $-30°$ Declination, the magnitudes of Pickering's "Harvard Photometry" are used. The places of the stars are for 1900.

Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY

In the designation of clusters and nebulæ the letter M. stands for "Messier," who made the first catalogue of 103 such objects in 1784; *e.g.*, 97 M. designates No. 97 on that list. A few objects from Herschel's catalogue are denoted by H with a number following.

The student or teacher who possesses a telescope is strongly urged to get Webb's "Celestial Objects for Common Telescopes." It is an invaluable accessory. (Longmans, Green & Co., N. Y.)

THE CIRCUMPOLAR CONSTELLATIONS.

We begin our study of Uranography with the constellations which are *circumpolar* (*i.e.*, within 40° of the north pole), because these are always visible in the United States, and so can be depended on to furnish land (or rather *sky*) -marks to aid in identifying and tracing out the others.

5. Ursa Major, the Great Bear (Map I.). — Of these circumpolar constellations none is more easily recognizable than Ursa Major. Assuming the time of observation as about eight o'clock in the evening on Sept. 22d (*i.e.*, 20^h sidereal time), it will be found below the pole and to the west. Hold the map so that the VIII. is at the bottom, and it will be rightly placed for the time assumed.

The familiar Dipper is sloping downward in the northwest, composed of seven stars, all of about the second magnitude excepting δ (at the junction of the handle to the bowl), which is of the third. The stars α and β are known as the "Pointers," because the line drawn from β through α, and produced about 30°, passes very near the Pole-star.

The dimensions of the Dipper furnish a convenient scale of angular measure. From α to β is 5°; α to δ is 10°; β to γ, 8°; from α to η at the extremity of the Dipper-handle (which is also the Bear's *tail*) is 26°.

6. The Dipper (known also in England as the "Plough," and as the "Wain," or wagon) comprises but a small part of

Digitized by Google

the whole constellation. The head of the Bear, indicated by a scattered group of small stars, is nearly on the line from β through α, carried on about 15°; at the time assumed (20^h sid. time), it is almost exactly below the pole. Three of the four paws are marked each by a pair of third or fourth magnitude stars $1\frac{1}{2}°$ or 2° apart. The three pairs are nearly equidistant, about 20° apart, and almost on a straight line parallel to the diagonal of the Dipper-bowl from α to γ, but some 20° south of it. Just now (20^h sid. time) they are all three very near the horizon for an observer in latitude 40°, but during the spring and summer they can be easily made out.

7. Names[1] of Principal Stars. —

α. DUBHE.	ϵ. ALIOTH.
β. MERAK.	ζ. MIZAR. The little star near it is *Alcor*, the "rider on his horse."
γ. *Phecda.*	η. BENETNASCH or *Alkaid.*
δ. *Megrez.*	

Double Stars: (1) ζ (Mizar), Mags. 3 and 5; Pos.[2] 149°; Dist. 14″.5. In looking at this object the tyro will be apt to think that the small star shown by the telescope is identical with Alcor: a very low power eye-piece will correct the error. (Astr. Fig. 113.) The large star is itself a "spectroscopic binary" (see Art. 465*). (2) ξ, the southern one of the pair which marks the left hind paw. *Binary:* Mags. 4 and 5; Pos. (1910) (about) 120°, Dist. (about) 3″. Position and distance both change rapidly, the period being only 61 years. This was the first binary whose orbit was computed.

Clusters and Nebulæ: (1) 81 and 82 M., A.R. $9^h\,45^m$, Dec. 69° 44′. Two nebulæ, one pretty bright, about half a degree apart. (2) 97 M., A.R. $11^h\,07^m$, Dec. 55° 43′—2° south-following β. A planetary nebula.

[1] Capitals denote names that are *generally* used; the others are met with only rarely.

[2] The "position angle" of a double star is the angle which the line drawn from the larger star to the smaller one makes with the hour-circle. It is always reckoned from the north completely around through the east, as shown in Fig. A.

Digitized by Google

3. Ursa Minor, the Lesser Bear (Map I.). — The line of the "pointers" unmistakably marks out the Pole-star ("Polaris" "Cynosura"), a star of the second magnitude standing alone. is at the end of the tail of Ursa Minor, or at the extremity the handle of the "*Little* Dipper," for in Ursa Minor, also, e seven principal stars form a dipper, though with the handle nt in a different ay from that of e other Dipper. ginning at "Po- ris" a curved line oncave towards rsa Major) drawn rough δ and ϵ ings us to ζ, where e handle joins the owl. Two bright ars (second and ird magnitude), β nd γ, correspond to e pointers in the rger Dipper, and e known as the Guardians of the ole": β is called *Kochab*." The remaining corner of the bowl is marked by ie faint star η with another still smaller one near it.



FIG. A. — Measurement of Distance and Position-Angle of a Double Star.

The Pole lies about $1\frac{1}{4}°$ from the Pole-star, on the line joining to ζ Ursæ Majoris (at the bend in the handle of the large ipper).

Telescopic Object. Polaris has a companion of the $9\frac{1}{2}$ magnitude, istant 18".6, — visible with a two-inch telescope.

9. Cassiopeia (Map I.). — This constellation lies on the pposite side of the pole from the Dipper at about the same

Digitized by Google

distance as the "Pointers," and is easily recognized by the zigzag, "rail-fence" configuration of the five or six bright stars that mark it. With the help of the rather inconspicuous star κ, one can make out of them a pretty good *chair* with the feet turned away from the pole. But this is wrong. In the recognized figures of the constellation the lady sits with feet *towards* the pole, and the bright star α is in her bosom, while ζ and the other faint stars south of α, are in her head and uplifted arms: ι, on the line from δ to ϵ produced, is in the foot. The order of the principal stars is easily remembered by the word *Bagdei*; *i.e.*, $\beta, \alpha, \gamma, \delta, \epsilon, \iota$.

Names of Stars: α (which is slightly variable) is known as SCHEDIR; β is called CAPH.

Double Stars: (1) η, Mags. 4–7½. Large star orange; small one purple. Pos. 170° ±, Dist. 5″.5. Binary, with a period of some 200 years. Easily recognized by its position about half-way between α and γ, a little off the line. (2) ψ, A.R. 1ʰ 17ᵐ, Dec. 67° 21′; Triple; Mags. 4½, 9 and 9; Pos. A to (B+C) 106°, Dist. 29″; Pos. B–C 257°, Dist. 2″.9. Found on a line from η through γ produced three times the distance η–γ: rather difficult for a four-inch telescope.

10. The Sidereal Time determined by the Apparent Position of Cassiopeia. — The line from the Pole-star through Caph or β Cassiopeiæ (which is the *leader* of all the bright stars of the constellation in their daily motion) is almost exactly parallel to the Equinoctial Colure. When, therefore, this star is vertically *above* the Pole-star **it is sidereal noon; it is 6ʰ when it is on the *great circle*** (not the parallel of altitude) drawn from the Pole-star to the west point of the Horizon; 12ʰ when vertically below it; and 18ʰ when due east. A little practice will enable one to read the sidereal time from this celestial clock with an error not exceeding 15 or 20 minutes.

11. Cepheus (Map I.). — This constellation contains very few bright stars. At the assumed time (20ʰ sidereal) it is above and west of Cassiopeia, not having quite reached the meridian above the pole. A line carried from α Cassiopeiæ through β, and produced 20° (distance $\alpha \ldots \beta = 5°$ nearly)

Digitized by Google

will pass very near to α Cephei, a star of the third magnitude, in the king's right shoulder. β Cephei is about 8° due north of α, and γ about 12° from β, both also of third magnitude: γ is so placed that it is at the obtuse angle of a rather flat isosceles triangle of which β Cephei and the Pole-star form the two other corners. Cepheus is represented as sitting behind Cassiopeia (his wife) with his feet upon the tail of the Little Bear, γ being in his left knee. His head is marked by a little triangle of fourth magnitude stars, δ, ε, and ζ, of which δ is a remarkable variable with a period of 5½ days (see Astr. Table IV.). There are several other small variables in the same neighborhood, but none of them are shown on the map.

Names of Stars: α is *Alderamin*, and β is *Alphirk*.

Double Stars: (1) β, Mags. 3 and 8; Pos. 251°; Dist. 14″. (2) δ, Mags. larger star 3.7 to 5 (variable), smaller one 7; Pos. 192°, Dist. 41″; Colors, yellow and blue. (3) κ, A.R. 20ʰ 13ᵐ, Dec. 77° 19′; Mags. 4.5 and 8.5; Pos. 124°; Dist. 7.″5; Colors, yellow and blue.

12. Draco (Map I.).—The constellation of Draco is characterized by a long, sinuous line of stars, mostly small, extending half-way around the pole and separating the two Bears. A line from δ Cassiopeiæ drawn through β Cephei and extended about as far again will fall upon the head of Draco, marked by an irregular quadrilateral of stars, two of which are of the 2½ and 3d magnitude. These two bright stars about 4° apart are β and γ; the latter in its daily revolution passes almost exactly through the zenith of Greenwich, and it was by observations upon it that the aberration of light was discovered (Astr. 125). The nose of Draco is marked by a smaller star, μ, some 5° beyond β, nearly on the line drawn through it from γ. From γ we trace the neck of Draco, eastward and downward[1] towards the Pole-star until we come to δ and ε and some smaller stars near them. There the direction of the line is reversed,

[1] The description here applies strictly only at 20ʰ sid. time.

Digitized by Google

so that the body of the monster lies between its own head and the bowl of the Little Dipper, and winds around this bowl until the tip of the creature's tail is reached at the middle of the line between the Pointers and the Pole-star. The constellation covers more than 12^h of right ascension.

13. One star deserves special notice: α, a star of the $3\frac{1}{2}$ magnitude which lies half-way between Mizar (ζ Urs. Maj.) and the Guards (β and γ Urs. Min.); 4700 years ago it was the Pole-star, within 10′ or 15′ of the pole, and much nearer than Polaris is at present, or ever will be. It is probable that its brightness has considerably diminished within the last 200 years; since among the ancient and mediæval astronomers it was always reckoned of the second magnitude.

Names of Stars: α is THUBAN; β, *Alwaid;* and γ, *Etanin.*

Double Stars: (1) μ, Mags. 4 and $4\frac{1}{2}$; Pos. 165°; Dist. 2″.5. Binary, with a probable period of about 600 years. (2) ϵ, Mags. 4, 8; Pos. 0°.0; Dist. 2″.9; yellow and blue. *Nebula*, A.R. $17^h\ 59^m$; Dec. 66° 38′. Planetary, like a star out of focus. This object is almost exactly at the pole of the ecliptic, about midway between δ and ζ Draconis, but a little nearer ζ.

14. Camelopardalis (Map I.).—This is the only remaining one of the strictly circumpolar constellations — a modern asterism containing no stars above fourth magnitude, and constituted by Hevelius simply to cover the great empty space between Cassiopeia and Perseus on one side, and Ursa Major and Draco on the other. The animal stands on the head and shoulders of Auriga, and his head is between the Pole-star and the tip of the tail of Draco.

The two constellations of Perseus (which at 20^h sidereal time is some 20° below Cassiopeia) and of Auriga are partly circumpolar, but on the whole can be more conveniently treated in connection with the equatorial maps. Capella, the brightest star of Auriga, and next to Vega and Arcturus the brightest star in the northern hemisphere, at the time assumed (Sept. 22, 8 P.M.), is a few degrees above the horizon in the N.E. Between it and the nose of Ursa Major is part of the constellation of the Lynx, — a modern asterism made, like Camelopardalis, merely to fill a gap.

Digitized by Google

15. The Milky Way in the Circumpolar Region. — The only circumpolar constellations traversed by it are Cassiopeia and Cepheus. It enters the circumpolar region from the constellation of Cygnus, which at 20^h sidereal time is just in the zenith, sweeps down across the head and shoulders of Cepheus, and on through Cassiopeia and Perseus to the northeastern horizon in Auriga. There is one very bright patch a degree or two north of β Cassiopeiæ; and half-way between Cassiopeia and Perseus there is another bright cloud in which is the famous cluster of the "Sword Handle of Perseus" — a beautiful object for even the smallest telescope.

---

16. Andromeda (Map II.). — Passing now to the equatorial maps and beginning with the northwestern corner of Map No. II., we come first to the constellation of Andromeda, which will be found exactly overhead in our latitudes about 10 o'clock in the middle of November, or at 8 o'clock a month later. Its characteristic configuration is the line of three second-magnitude stars, α, β, and γ, extending east and north from α, which itself forms the N.E. corner of the so-called "Great Square of Pegasus," and is sometimes lettered as δ Pegasi. This star may readily be found by extending an imaginary line from Polaris through β Cassiopeiæ, and producing it about as far again: α is in the head of Andromeda, β in her waist, and γ in the left foot. About half-way from α to β, a little south of the line, is δ (of the third magnitude) with π and ϵ of the fourth magnitude near it. A line drawn northwesterly from β nearly at right angles to the line $\beta\gamma$, will pass through μ at a distance of about 5°, and produced another 5° will strike the "great nebula" (Astr. 470), which forms a little obtuse-angled triangle with ν and a sixth-magnitude star known as 32 Andromedæ.

Andromeda has her mother, Cassiopeia, close by on the north, and at her feet is Perseus, her deliverer, while her head rests upon the

Digitized by Google

shoulder of Pegasus, the winged horse which brought Perseus to her rescue. To the south, beyond the intervening constellations of Aries and Pisces, Cetus, the sea-monster, who was to have devoured her, stretches his ungainly bulk.

Names of Stars. α, *Alpheratz*; β, *Mirach*; γ, *Almaach*.

Double Stars. (1) γ, Mags. 3, 5; Pos. 62°; Dist. 11″; colors, orange and greenish blue — a beautiful object. The small star is itself double, but at present so close as to be beyond the reach of any but very large instruments (Astr. Fig. 113). (2) π (2° N. and a little west of δ), Mags. 4, 9; Pos. 174°; Dist. 36″; white and blue.

Nebulæ. M. 31; the great nebula; visible to naked eye. M. 32; small, round, and bright, is in the same low-power field with 31; south and east of it.

17. Pisces (Map II.). — Immediately south of Andromeda lies Pisces, the first of the zodiacal constellations, though now occupying (in consequence of precession) the *sign* of Aries. It has not a single conspicuous star, and is notable only as containing the vernal equinox, or first of Aries, which lies near the southern boundary of the constellation in a peculiarly starless region. A line from α Andromedæ through γ Pegasi continued as far again strikes about 2° east of the equinox.

The body of the southern fish lies about 15° south of the middle of the southern side of the great square of Pegasus, and is marked by an irregular polygon of small stars, 5° or 6° in diameter. A long crooked "ribbon" of little stars runs eastward for more than 30°, terminating in α Piscium, called *El Rischa*, a star of the fourth magnitude 20° south of the head of Aries. From there another line of stars leads up N.W. in the direction of δ Andromedæ to the northern fish, which lies in the vacant space south of β Andromedæ.

Double Stars. (1) α, Mags. 4, 5.5; Pos. 324°; Dist. 3″. (2) ψ' (2° S.E. of η Andromedæ — see map), Mags. 4.9, 5; Pos. 160°; Dist. 31″.

18. Triangulum (Map II.). — This little constellation, insignificant as it is, is one of Ptolemy's ancient 48. It lies half-way between γ Andromedæ and the head of Aries, characterized by three stars of the third and fourth magnitudes.

Digitized by Google

Double Stars. (1) ι or 6 (5° nearly due south of β Trianguli, and at the obtuse angle of an isosceles triangle of which α and γ are the other two corners), Mags. 5, 6.5; Pos. 76°; Dist. 4″; topaz-yellow and green.

19. Aries (Map II.). — This is the second of the zodiacal constellations (now occupying the *sign* of Taurus). It is bounded north by Triangulum and Perseus, west by Pisces, south by Cetus, and east by Taurus. The characteristic star-group is that composed of α, β, γ (see map), about 20° due south of γ Andromedæ: α, a star of the 2½ magnitude is fairly conspicuous, forming as it does a large isosceles triangle with β and γ Andromedæ.

Names of Stars. α, *Hamal*; β, *Sheratan*; γ, *Mesartim*.

Double Stars. (1) γ, Mags. 4.5, 5; Pos. 0°; Dist. 8″.8. (This is probably the earliest known double star; noticed by Hooke in 1664.) (2) ε, Mags. 5, 6.5; Pos. 200°; Dist. 1″.2. (About one-third of the way from α Arietis towards Aldebaran, ζ is 4° beyond it on the same line.) This is probably too difficult for any instrument less than 4 or 4½ inches' aperture. (3) π, Triple; Mags. 5, 8.5, and 11; A–B, Pos. 122°; Dist. 3″.1; A–C, Pos. 110°; Dist. 25″. (At the southern corner of a nearly isosceles triangle formed with ε and τ, ε being at the obtuse angle.)

The star 41 Arietis (3½ mag.), which forms a nearly equilateral triangle with α Arietis and γ Trianguli, constitutes, with two or three other small stars near it, the constellation of *Musca* (Borealis), a constellation, however, not now generally recognized.

20. Cetus (Map II.). — South of Aries and Pisces lies the huge constellation of Cetus, which backs up into the sky from the southeastern horizon. The head lies some 20° S.E. of α Arietis, marked by an irregular pentagon of stars, each side of which is 5° or 6° long. The southern edge of it is formed by the stars α (2½ mag.) and γ (3½ mag.): δ lies nearly south of γ. β, the brightest star of the constellation (2d magnitude), stands alone nearly 40° west and south of α. About half-way

Digitized by Google

from β to γ the line joining them passes through a characteristic quadrilateral (see map), the N.E. corner of which is composed of two fourth-magnitude stars, ζ and χ. The remarkable variable o Ceti (Mira) lies almost exactly on the line joining γ and ζ, a little nearer to γ than to ζ. It is visible to the naked eye for about a month or six weeks every eleven months, when near its maximum.

Names of Stars. α, MENKAR; β, *Diphda* or *Deneb Kaitos*; ζ, *Baten Kaitos*; o, MIRA.

Double Stars. (1) γ, Mags. 3.5, 7; Pos. 290°; Dist. 2″.5; yellow and blue.

South of Cetus lies the constellation of Sculptoris Apparatus (usually known simply as Sculptor), which, however, contains nothing that requires notice here. South of Sculptor, and close to the horizon, even when on the meridian, is Phœnix. It has some bright stars, but none easily observable in the United States.

21. Perseus (Maps I. and II.). — Returning now to the northern limit of the map, we come to the constellation of Perseus. Its principal star is α, rather brighter than the standard second magnitude, situated very nearly on the prolongation of the line of the three chief stars of Andromeda. A very characteristic configuration is "the segment of Perseus" (Map I.), a curved line, formed by δ, α, γ, and η, with some smaller stars, concave towards the northeast, and running along the line of the Milky Way towards Cassiopeia. The remarkable variable star β, or Algol (Astr. 453), is situated about 9° south and a little west of α, at the right angle of a right-angled triangle which it forms with α (Persei) and γ Andromedæ. Some 8° south and slightly east of δ is ϵ, and 8° south of ϵ are ζ and o of the fourth magnitude in the foot of the hero. Algol and a few small stars near it form "Medusa's Head."

Names of Stars. α is *Marfak*, or ALGENIB; β is ALGOL.

Double Stars. (1) ϵ, Mags. 3.5, 9; Pos. 10°; Dist. 8″.4. (2) ζ

Digitized by Google

Quadruple; Mags. 3.5, 10, 11, 12; Pos. A–B, 207°; Dist. 13″.2, 83″, 121″. (3) η, Mags. 5, 8.5; Pos. 300°; Dist. 28″; orange and blue.

Clusters. (1) ♅ VI. 33 and 34. Magnificent. Half-way between γ Persei and δ Cassiopeiæ. (2) M. 34; A.R. $2^h\ 34^m$; Dec. 42° 11′; coarse, with a pretty double star (eighth mag.) included.

22. Aurīga (Maps I. and II.). — Proceeding east from Perseus we come to Auriga, instantly recognized by the bright yellow star CAPELLA (the Goat) and her attendant Hœdi (or Kids). Capella, α Aurigæ, according to Pickering, is precisely of the same brightness as Vega (Mag. = 0.2), both of them being about $\frac{1}{5}$ of a magnitude fainter than Arcturus, but distinctly brighter than any other stars visible in our latitudes except Sirius itself. About 10° east of Capella is β Aurigæ of the second magnitude, and 8° south of β is θ of the third magnitude; δ Aurigæ is 10° north of β in the circumpolar region. ϵ, ζ, and η, 4° or 5° S.W. of α, are the "Kids."

Names of Stars. α, CAPELLA; β, MENKALINAN.

Double Stars. (1) ω, Mags. 5, 9; Pos. 353°; Dist. 7″; white, light blue. β is a spectroscopic double (see Art. 465*).

Clusters. (1) M. 37; A.R. $5^h\ 44^m$; Dec. 32° 31′ (on the line from θ Aurigæ to ζ Tauri, one-third of the way from θ). Fine for small instrument. (2) M. 38; A.R. $5^h\ 21^m$; Dec. 35° 47′. Nearly at the middle of the line from θ to ω. (3) M. 36; A.R. $5^h\ 28^m$; Dec. 34° 3′. One-third of the way from M. 38 to M. 37.

23. Taurus (Map II.). — This, the third of the zodiacal constellations, is bounded north by Perseus and Auriga, west by Aries, south by Eridanus and Orion, and east by Orion and Gemini. It is unmistakably characterized by the Pleiades, and by the V-shaped group of the Hyades which forms the face of the bull, with the red ALDEBARAN (α Tauri) blazing in the creature's eye, as he charges down upon Orion. His horns reach out towards Gemini and Auriga, and are tipped with the second and third magnitude stars β and ζ. As in the case of Pegasus, only the head and shoulders appear in the constella-

Digitized by Google

tion. Six of the Pleiades are easily visible, and on a dark night a fairly good eye will count nine (see Astr. 469). With a 3-inch telescope about 100 stars are visible in the cluster. In the Hyades the pretty naked-eye double θ_1, θ_2, is worth noting.

Names of Stars. α, ALDEBARAN; β, *El Nath*; η (the brightest of the Pleiades), ALCYONE. For the names of the other Pleiades, see the figure in Art. 469 of the Astronomy.

Double Stars. (1) α has a small, distant companion, 12th magnitude; Pos. 36°; Dist. 1′ 48″. It has also a second companion much nearer and more minute, but far beyond the reach of ordinary telescopes. (2) τ, Mags. 5 and 8; Pos. 210°; Dist. 62″; white and violet. Found by drawing a line from γ (at the point of the V of the Hyades) through ϵ, and producing it as far again.

Nebula. M. 1; A.R. $5^h\ 27^m$; Dec. 21° 56′, about 1° west and a little north of ζ. Often mistaken for a comet. The so-called "Crab Nebula."

24. Orion (not O′-rĭ-on) (Map II.). — On the whole this is the finest constellation in the heavens. As he stands facing the bull his shoulders are marked by the two bright stars, α and γ, the former of which in color and brightness closely matches Aldebaran. In his left hand he holds up the lion skin, indicated by the curved line of little stars between γ and the Hyades. The top of the club, which he brandishes in his right hand, lies between ζ Tauri and μ and η Geminorum. His head is marked by a little triangle of stars of which λ is the chief. His belt consists of three stars of the second magnitude pointing obliquely downward towards Sirius. It is very nearly 3° in length, with the stars in it equidistant like a measuring-rod, so that it is known in England as the "Ell and Yard." From the belt hangs the sword, composed of three smaller stars lying more nearly north and south: the middle one of them is the multiple θ in the great nebula. β Orionis, or RIGEL, a magnificent white star, is in the left foot, and κ is in the right knee. Orion has no right *foot*, or if he has, it is

Digitized by Google

hidden behind Lepus. The quadrilateral α, γ, δ, κ, with the diagonal belt δ, ϵ, ζ, once learned can never be mistaken for anything else in the heavens.

25. *Names of Stars.* α, BETELGEUSE; β, RIGEL; γ, *Bellatrix*; κ, *Saiph*; δ, *Mintaka*; ϵ, *Alnilam*; ζ, *Alnitak*.

Double Stars. In these Orion is remarkably rich. (1) β (Rigel), Mags. 1 and 9; Pos. 200°; Dist. 9″.5; both white, — a beautiful and easy object. (2) δ (the westernmost star in the belt), Mags. 2.5 and 7; Pos. 0; Dist. 53′. (3) ζ, Triple; Mags. 2.5, 6.5, 10; A–B, Pos. 155°, Dist. 2″.4; A–C, Pos. 9°, Dist. 59″. (4) ι, Triple; Mags. 3.5, 8.5, 11; A–B, Pos. 142°, Dist. 11″.5; A–C, Pos. 103°, Dist. 49″. (This is the lowest star in the sword, just below the nebula.) (5) θ, Multiple, the trapezium in the nebula. Four stars are easily seen by small telescopes (Astr. Fig. 113). (6) σ, Triple; Mags. 4, 8, 7; A–B, Pos. 84°, Dist. 12″.5; A–C, Pos. 61°, Dist. 42″ (1½° S.W. of ζ).

Nebula. M. 42; attached to the multiple star θ. *The* nebula of all the heavens; by far the finest known, though in a small telescope wanting much of the beauty brought out by a larger one.

26. Eridănus (Map II.). — This constellation lies south of Taurus, in the space between Cetus and Orion, and extends far below the southern horizon. Its brightest star α (ACHERNAR) is never visible in the United States.

Starting with β of the third magnitude, about 3° north and a little west of Rigel (β Orionis), one can follow a sinuous line of stars, some of them of the third and fourth magnitudes, westward about 30° to the paws of Cetus, 10° south of α Ceti; there the stream turns at right angles southwards for 10°, then southeast for about 20°, and finally southwestward to the horizon. One could succeed in fully tracing it out, however, only by help of a map on a larger scale than the one we are able to present.

Names of Stars. β, *Cursa*; γ, *Zaurack*.

Double Stars. (1) 32; A.R. $3^h\ 48^m$, Dec. S. 3° 19′; Mags. 5, 7; Pos. 347°; Dist. 6″.6; yellow, blue; very fine. (2) o^2, Triple; A.R.

Digitized by Google

$4^h\ 10^m$; Dec. S. 7° 50′; Mags. 5, 10, 10; Pos. A, $\frac{(B+C)}{2}$, 108°; Dist. 83″; Pos. B–C, 110°, Dist. 4″; very pretty.

27. Lepus (Map II.). — This little constellation (one of the ancient 48) lies just south of Orion, occupying a space of some 15° square. Its characteristic configuration is a quadrilateral of third and fourth magnitude stars, with sides from 3° to 5° long, 10° south of κ Orionis, and 15° west and a little south of Sirius.

Double Stars. (1) γ (the S.E. corner of the quadrilateral) is a coarse double. Mags. 4, 6.5; Pos. 350°; Dist. 93″. (2) κ ($5\frac{1}{2}$° south of Rigel), Mags. 5 and 9; Pos. 0°; Dist. 3″.*i*.

28. Columba (Noah's Dove) (Map II.). — This is next south of Lepus: too far south to be well seen in the Northern States. Its principal star, α, or *Phact*, of the $2\frac{1}{2}$ magnitude, is readily found by drawing a line from Procyon to Sirius, and prolonging it nearly the same distance. And in passing we may note that a similar line drawn from α Orionis through Sirius and produced, will strike near ζ Argus, or "*Naos*," a star about as bright as Phact, — the two lines which intersect at Sirius making the so-called "*Egyptian X*."

29. Lynx (Map I., II., and III.). — Returning now to the northern limit of the map, we find the modern constellation of the Lynx lying just east of Auriga and enveloping it on the north and in the circumpolar region. It contains no stars above the fourth magnitude, and is of no importance except as occupying an otherwise vacant space.

Double Stars. (1) 38, or ρ Lyncis, A.R. $9^h\ 11^m$; Dec. 37° 21′; Mags. 4, 7.5; Pos. 240°; Dist. 2″.9; white and lilac. (This is the northern one of a pair of stars which closely resembles the three pairs that mark the paws of Ursa Major. This pair makes nearly an isosceles triangle with the two pairs $\lambda\ \mu$ and $\iota\ \kappa$, Ursæ Majoris — see map.)

30. Gemini (Map II.). — This is the fourth of the zodiacal constellations (mostly in the *sign* of Cancer), containing the summer solstitial point about 2° west and a little north of the star η. It lies northeast of Orion and southeast of Auriga, and is sufficiently characterized by the two stars α and β (about $4\frac{1}{2}$° apart), which mark the heads of the twins. The southern

Digitized by Google

one, β, or POLLUX, is the brighter, but α (CASTOR) is much the more interesting, as being double. The feet are marked by the third-magnitude stars γ and μ, some 10° east of ζ Tauri, and the map shows how the lines that join these to β and α respectively mark the places of δ and ϵ. η, 2° west of μ, is a variable, and is also double, though as such it is beyond the power of ordinary telescopes.

Names. α, CASTOR; β, POLLUX; γ, *Alhena*, μ, *Tejat (Post)*; η, *Tejat (Prior)*; δ, *Wasat*; ϵ, *Meboula*.

Double Stars. (1) α, Mags. 2.5, 3; Pos. 225°; Dist. 5″.5. Binary; period undetermined, but certainly over 200 years. The larger of the close pair is also a spectroscopic binary, with period of about 3 days (see Art. 465*). There is also a companion of ninth mag., distant about 74″; Pos. 164°. (2) δ, Mags. 3, 8; Pos. 203°; Dist. 7″. (3) μ, Mags. 3, 11; Pos. 79°; Dist. 80″.

Nebulæ and Clusters. (1) M. 35; A.R. $6^h\ 01^m$; Dec. 24° 21′; N.W. of η at the same distance as that from μ to η, and on the line from γ through η produced. The map is not quite right in this respect. (2) ♅ IV. 45; A.R. $7^h\ 22^m$; N. 21° 10′. A nebulous star in a small telescope; in a large telescope, very peculiar — 2° southeast of δ.

31. Canis Minor (Map II.). — This constellation, just south of Gemini, is sufficiently characterized by the bright star PROCYON, which is 25° due south of the mid-point between Castor and Pollux. α, β, and γ together form a configuration closely resembling that formed by α, β, and γ Arietis. Procyon, α Orionis, and Sirius form nearly an equilateral triangle with sides of about 25°.

Names. α, PROCYON; β, *Gomelza*.

Double Stars. (1) Procyon has a small companion, Dist. 40″, Pos. 312°, — too small, however, for anything less than an 8-inch telescope. In 1896 a still smaller companion, like that of Sirius, was found much nearer the large star. (See Art. 464.) (2) (Σ 1126) (following Procyon 43^s, and 2′ south, — the brightest of the stars in that field), Mags. 7, 7.5; Pos. 145°; Dist. 1″.5; a good test for a 4-inch glass.

Digitized by Google

32. Monoceros (The Unicorn) (Map II.). — This is one of the ew constellations organized by Hevelius to fill the gap between Gem- ıi and Canis Minor on the north, and Argo Navis and Canis Major n the south. It lies just east of Orion. It has no conspicuous stars, ut is traversed by a brilliant portion of the Milky Way. The α fourth mag.) of the constellation lies about half-way between α rionis and Sirius, a little west of the line joining them.

Double Stars. (1) 8, or *b* (7½° east and 3° south of α Orionis), lags. 5, 8; Pos. 24°; Dist. 12″.9; colors, orange and lilac. A fine ow-power field. (2) 11 Monocerōtis, a fine *triple* (see Fig. 113 of str.), A.R. 6ʰ 24ᵐ; Dec. *south* 6° 57′; A to B–C, Pos. 130°, Dist. 8″; –C, Pos. 120°, Dist. 2″.5. The star is very nearly pointed at by a ne drawn from ζ Canis Majoris, north through β, and continued as ar again.

Clusters. (1) M VII. 2; A.R. 6ʰ 24ᵐ; Dec. N. 5° 2′ (visible to the aked eye about 1½° N.E. of 8 Monocerotis described above). A fine luster for a low power. (2) M. 50; A.R. 6ʰ 57ᵐ; Dec. S. 8° 9′. n the Milky Way, on the line from Sirius to Procyon, two-fifths of ne distance.

33. Canis Major (Map II.). — This glorious constellation eeds no description. Its α is the Dog Star, SIRIUS, beyond ll comparison the brightest in the heavens, and probably one f our nearer neighbors. It is nearly pointed at by the line rawn through the three stars of Orion's belt. β, at the ex- remity of the uplifted paw, is of the second magnitude, and o are several of those farther south in the rump and tail of he animal, who sits up watching his master Orion, but with n eye out for Lepus.

Names. α, SIRIUS; β, *Mirzam*; γ, *Muliphen*; δ, *Wesen*; ε, *Adara*. is said to have disappeared from 1670 to 1690, but at present is not ecognized as variable, though much fainter than would be expected om its being lettered as γ.

Double Stars. (1) Sirius itself has a small companion (see Art. 63). (2) μ (4° N.E. of Sirius), Mags. 5, 9.5; Pos. 335°; Dist. 3″.5.

Clusters. (1) M. 41 (4° south of Sirius); a fine group with a red ar near centre.

Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY

34. Argo Navis (The Ship) (Maps II. and III.). — This is one of the largest, most important, and oldest of the constellations, lying south and east of Canis Major. Many Uranographers now divide it into *three*, Puppis, Vela, and Carina. Its brightest star, α Argûs, Canopus, ranks next to Sirius, but is not visible anywhere north of the parallel of 38°. The constellation, huge as it is, is only a *half* one, like Pegasus and Taurus, — only the stern of a vessel, with mast, sail, and oars; the stem being wanting. In the part of the constellation covered by our maps the most conspicuous stars lie east and southeast of Canis Major. We have already mentioned ζ, or *Naos* (Art. 28), at the southeast extremity of the "Egyptian X"; and about 8° south and a little east of it is γ, nearly of the second magnitude.

Clusters. One or two clusters are accessible in our latitudes. (1) (H VIII. 38), A.R. $7^h\ 31^m$; Dec. S. 14° 12′. Pointed at by the line from β Can. Maj. through Sirius, continued $2\frac{1}{2}$ times as far. Visible to naked eye; rather coarse. (2) M. 46; a little more than 1° east and south of the preceding. (3) M. 93, A.R. $7^h\ 39^m$; Dec. S. 23° 34′; about 2° N.W. of ζ Argûs.

35. Cancer (Maps II. and III.). — This is the fifth of the zodiacal constellations, bounded north by Lynx and Leo Minor, south by the head of Hydra, west by Gemini and Canis Minor, and east by Leo. It does not contain a single conspicuous star, but is easily recognizable from its position, and in a dark night by the nebulous cloud known as "Præsepe," or the "Manger," with the two stars γ and δ near it, — the so-called "Aselli," or "Donkeys." Præsepe (sometimes also called the "Beehive") is really a coarse cluster of seventh and eighth magnitude stars, resolvable by an opera-glass. The line from Castor through Pollux, produced about 12°, passes near enough to it to serve as a pointer. α, of the fourth magnitude, is on the line drawn from Præsepe through δ (the southern Asellus), produced about 7°; β may be recognized by drawing a line from γ (the northern Asellus) through Præsepe, and continuing it about 12°.

Double Stars. (1) ι, Mags. 4, 6.5; Pos. 308°; Dist. 30″; orange and blue; nearly due north of γ, distance twice that between the Aselli.

Digitized by Google

(2) ζ, Triple (see Astr. Fig. 113); A–B, Mags. 6 and 7, Pos. (1905) 346°, Dist. 1″; in rapid motion; period about 60 years. A–C, Pos. (1905) 110°, Dist. 5″; also in motion, but period unknown and much longer. Easily found by a line from α Gem. through β, produced two and a half times as far.

36. Leo (Map III.). — East of Cancer lies the noble constellation of Leo, which adorns the evening sky in March and April; it is the sixth of the zodiacal constellations, now occupying the sign of Virgo. Its leading star REGULUS, or "*Cor Leonis*," is of the first magnitude, and two others, β and γ, are of the second. α, γ, δ, and β form a conspicuous irregular quadrilateral (see map), the line from Regulus to Denebola being 26° long. Another characteristic configuration is "The Sickle," of which α, η is the handle, and the curved line η, γ, ζ, μ, and ε is the blade, the cutting edge being turned towards Cancer. The "radiant" of the November meteors lies between ζ and ε.

Names. α, REGULUS; β, DENEBOLA; γ, *Algeiba*; δ, *Zosma*.

Double Stars. (1) γ, Mags. 2, 3.5; Pos. 116°; Dist. 3″.4; binary; period about 400 years. (2) ι, Mags. 4 and 7; Pos. 65°; Dist. 2″.5; yellow and bluish; easily recognized by aid of the map. (3) 54, Mags. 4.5, 7; Pos. 103°; Dist. 6″.2. Found by producing the line from β through δ half its length.

37. Leo Minor and Sextans (Map III.). — *Leo Minor* is an insignificant modern constellation composed of a few small stars north of Leo, between it and the hind feet of Ursa Major. It contains nothing deserving special notice. A similar remark holds as to *Sextans* even more emphatically.

38. Hydra (Map III.). — This constellation, with its riders Crater and Corvus, is a large and important one, though not very brilliant. The head is marked by a group of five or six fourth and fifth magnitude stars just 15° south of Præsepe. A curving line of small stars leads down southeast to α, "*Cor Hydræ*," a small second or bright third magnitude star stand-

Digitized by Google

ing very much alone. From there, as the map shows, an irregular line of fourth-magnitude stars running far south and then east, almost to the boundary of Scorpio, marks the creature's body and tail, the whole covering almost six hours of right ascension, and very nearly 90° of the sky. About the middle of his length, and just below the hind feet of Leo (30° due south from Denebola), we find the little constellation of *Crater;* and just east of it the still smaller but much more conspicuous one of *Corvus*, with two second-magnitude stars in it, and four of the third and fourth magnitudes. It is well marked by a characteristic quadrilateral (see map), with δ and η together at its northeast corner. The order of the letters differs widely from that of brightness in this constellation, suggesting that changes may have occurred.

Names. α Hydræ, ALPHARD or *Cor Hydræ;* α Crateris, *Alkes*; α Corvi, *Alchiba;* δ Corvi, *Algores.*

Double Stars. (1) ε Hydræ (the northernmost one of the group that marks the head), Mags. 4, 8; Pos. 220°; Dist. 3″.5; yellow and purple. (2) δ Corvi, Mags. 3, 8; Pos. 210°; Dist. 24″; yellow and purple. (3) *Nebula,* ♅ IV. 27, A.R. $10^h\ 19^m$; Dec. S. 18° 2′ (3° S. and ½° W. of μ — see map). Bright planetary nebula, about as large as Jupiter.

39. Virgo (Map III.). — East and south of Leo lies Virgo, the seventh zodiacal constellation, bounded on the north by Boötes and Coma Berenices, on the east by Boötes and Libra, and on the south by Corvus and Hydra. Its α, SPICA Virginis, is of the 1½ magnitude and, standing rather alone 10° south of the celestial equator, is easily recognized as the southern apex of a nearly equilateral triangle which it forms with Denebola (β Leonis) to the northwest, and Arcturus northeast of it. β Virginis of the third magnitude is 14° due south of Denebola. A line drawn eastward and a little south from β (third magnitude) and then carried on, curving northward, passes successively (see map) through η, γ, δ, and ε,

Digitized by Google

of the third magnitude (notice the word formed by the letters *Bēgde*, like *Bagdei* in Cassiopeia, Art. 9). θ lies nearly midway between α and δ. There are also a number of other fourth-magnitude stars.

Names. α, SPICA and *Azimech*; β, *Zavijava*; ϵ, *Vindemiatrix*.

Double Stars. (1) γ, (Binary; period 185 years; not quite half-way from Spica to Denebola, and a little west of the line), Mags. 3, 3; Pos. (1905) 328°; Dist. 6″.2; very easy and fine (Astr. Fig. 113). (2) θ (two-fifths of the way from *Spica* towards δ), Triple; Mags. A 4.5, B 9, C 10; Pos. A–B, 345°, Dist. 7″; A–C, Pos. 295°, Dist. 65″. (3) o (one-third of the way from *Denebola* towards γ Virginis), Mags. 6 and 8; Pos. 228°; Dist. 3″.5. Spica is a spectroscopic binary (Art. 465*).

Nebulæ. (1) M. 49; A.R. $12^h\ 24^m$; Dec. $+8°\ 40'$. Forms an equilateral triangle with δ and ϵ. It lies in the remarkable "nebulous" region of Virgo. But most of the nebulæ are faint, and observable only with large telescopes. (2) ♅ II. 74 and 75; A.R. $12^h\ 47^m$; Dec. $+11°\ 53'$; two in one field, 2° west and a little north of ϵ. (3) M. 86 (midway between Denebola and ϵ); A.R. $12^h\ 20^m$; Dec. $+13°\ 36'$. A large telescope shows nearly a dozen nebulæ within 2° of this place.

40. Coma Berenices (Map III.). — This little constellation, composed of a great number of fifth and sixth magnitude stars, lies 30° north of γ and η Virginis, and about 15° northeast of Denebola. It contains a number of interesting double stars, but they are not easily found without the help of an equatorial mounting and graduated circles.

41. Canes Venatici (The Hunting Dogs). — These are the dogs with which Boötes is pursuing the Great Bear around the pole: the northern of the two is *Asterion*, the southern *Chara*. Most of the stars are small, but α is of the $2\frac{1}{2}$ magnitude, and is easily found by drawing from η Ursæ Majoris (the star in the end of the Dipper-handle) a line to the southwest, perpendicular to the line from η to ζ (Mizar) and about 15° long: it is about one-third of the way from η Ursæ Majoris to δ Leonis. With Arcturus and Denebola it forms a triangle much like that which they form with Spica.

Digitized by Google

Names. α is known as Cor Caroli (Charles II. of England).

Double Stars. (1) α, or 12 Canum, Mags. 3 and 5; Pos. 227°; Dist. 20″. (2) 2 Canum (one-third of the way from α towards δ Leonis), Mags. 6 and 8; Pos. 260°; Dist. 41″.3; orange, smalt blue.

Nebulæ. (1) M. 51; A.R. $13^h\ 25^m$; Dec. 47° 49′ (3° west and somewhat south of Benetnasch). A faint double nebula in small telescopes; in great ones, the wonderful "Whirlpool Nebula" of Lord Rosse. (2) M. 3; bright cluster (half a degree north of the line from α Canum to Arcturus, and a little nearer the latter). It is one of the *variable-star clusters* discovered in 1895 (see Art. 455*).

42. Boötes (Maps III. and I.). — This fine constellation is bounded on the west by Ursa Major, Canes Venatici, Coma Berenices, and Virgo, and on the south by Virgo. It extends more than 60° in declination, from near the equator quite to Draco, where the uplifted hand overlaps the tail of the Bear. Its principal star, ARCTURUS, is of a ruddy hue, and in brightness is excelled only by Sirius among the stars visible in our latitudes. Canopus and α Centauri are reckoned brighter, but they are southern circumpolars. Arcturus is at once recognized by its forming with Spica and Denebola the great triangle already mentioned (Art. 39). Six degrees west and a little south of it is η, of the third magnitude, which forms with it, in connection with ν, a configuration like that in the head of Aries. ϵ is about 10° northeast of Arcturus, and in the same direction about 10° farther lies δ. A pentagon is formed by these two stars along with β, γ, and ρ. "Boötes" means "the shouter" (or, according to others, "the herdsman").

Names. α, ARCTURUS; β, *Nekkar;* ϵ, *Izar;* η, *Muphrid*, γ, *Seginus.*

Double Stars. (1) ϵ, Mags. 3, 6; Pos. 325°; Dist. 3″.1; orange and greenish blue; very fine. (2) ζ (about 9° southeast from Arcturus, at right angles to the line $\alpha\epsilon$), Mags. 3.5, 4; Pos. 295°; Dist. 0″.8; a good test for a 4-inch glass. (3) π ($2\frac{1}{2}$° north of ζ), Mags. 4.9, 6; Pos. 101°; Dist. 5″.3. (4) ξ (10° due east from Arcturus, 3° N.E. from π), Mags. 4.7, 6.6; Pos. (1905) 176°; Dist. 2″.5; yellow and purple. Binary; period 127 years.

Digitized by Google

43. Corona Borealis (Map III.). — This beautiful little constellation lies 20° northeast of Arcturus, and is at once recognizable as an almost perfect semicircle composed of half a dozen stars, among which the brightest, α, is of the second magnitude. The extreme northern one is θ; next comes β, and the rest follow in the $\beta\ \alpha\ \gamma\ \delta\ \epsilon\ \iota$ (*Bagdei*) order, just as in Cassiopeia.

Names. α, *Gemma*, or *Alphacca*.

Double Stars. (1) ζ (nearly pointed at by ϵ–δ Boötis; 7° from ϵ), Mags. 5, 6; Pos. 301°; Dist. 6″; white and greenish. (2) η, rapid binary, at certain times can be split by a 4-inch glass. Mags. 6, 6.5; pointed at by the line from α through β, 2° beyond β. The temporary star of 1866 (Astr. 450) lies 1½° S.E. of ϵ Coronæ.

44. Libra (Map III.). — This is the eighth of the zodiacal constellations, and lies east of Virgo, bounded on the south by Centaurus and Lupus, on the east by the upstretched claw of Scorpio, and on the north by Serpens and Virgo. It is inconspicuous, the most characteristic figure being the trapezoid formed by the lines joining the four stars α, ι, γ, β. β, which is the northernmost of the four, is the brightest (2½ magnitude), and is about 30° nearly due east from Spica, while α is about 10° southwest of β. The remarkable variable δ Libræ is 4° west and a little north from β. Most of the time it is of the 4½ or fifth magnitude, but runs down nearly two magnitudes at the minimum.

Names. α, *Zuben el Genubi*; β, *Zuben el Chamali*.

Cluster. M. 5; A.R. $15^h\ 12^m$; Dec. N. 2° 32′. This is within the boundaries of *Serpens*, and just a little north and west of the fifth-magnitude star 5 Serpentis. It is a *variable-star cluster* (Art. 455*).

45. Antlia, Centaurus, and Lupus (Map III.). — These constellations lie south of Hydra and Libra. *Antlia Pneumatica* (the "Air-Pump") is a modern constellation of no importance and hardly recognizable by the eye, having only a single star as bright as the 4½ mag.

Digitized by Google

nitude. *Centaurus*, on the other hand, is an ancient and extensive asterism, containing in its (south) circumpolar portion two stars of the first magnitude: α Centauri stands next after Sirius and Canopus in brightness, and, as far as present knowledge indicates, is our nearest neighbor among the stars. The part of the constellation which becomes visible in our latitudes is not specially brilliant, though it contains several stars of the $2\frac{1}{2}$ and third magnitude in the region that lies south of Corvus and Spica Virginis. A line from ϵ Virginis through Spica, produced a little more than its own length, will strike very near θ, a solitary star of the $2\frac{1}{2}$ magnitude in the Centaur's left shoulder. ι (third mag.) lies 11° west of θ, and η (third mag.) 9° southeast; while 5° or 6° south of the line from θ to ι lies a tangle of third-magnitude stars, which, if they were at a higher elevation, would be conspicuous. Centaurus is best seen in May or early in June.

Lupus, also one of Ptolemy's constellations, lies due east of Centaurus and just south of Libra. It contains a considerable number of third and fourth magnitude stars; but is too low for any satisfactory study in our own latitudes. It is best seen late in June. These constellations contain numerous objects interesting for a southern observer, but nothing available for our purpose.

46. Scorpio (*or Scorpius*) (Map III.). — This, the ninth of the zodiacal constellations, and the most brilliant of them, lies southeast of Libra, which in ancient times used to form its claws (*Chelæ*). It is bounded north by Ophiuchus, south by Lupus, Norma, and Ara, and east by Sagittarius. It is recognizable at once on a summer evening by the peculiar configuration, like a boy's kite, with a long streaming tail reaching far down to the southern horizon. Its principal star, ANTARES, is of the first magnitude and fiery red, like the planet Mars. From this it gets its name, which means "the rival of *Ares*" (*Mars*). β (second magnitude) is in the arch of the kite bow, about 8° or 9° northwest of Antares, while the star which Bayer lettered as γ Scorpii is well within Libra, 20° west of Antares. (There is no little discordance and confusion among Uranographers as to the boundary between the

Digitized by Google

two constellations.) The other principal stars of the constellation are easily found on the map; δ is 3° southwest of β, while ϵ, ζ, η, θ, ι, κ, and λ follow along in order in the tail of the creature, except that between ϵ and ζ is interposed the double μ. ϵ, θ, and λ are all of the second magnitude, and the others of the third.

47. *Names.* α, ANTARES; β, *Akrab.*

Double Stars. (1) α, Mags. 1 and 7; Pos. 270°; Dist. 3.″5; fiery red and vivid green. A beautiful object when the state of the air allows it to be well seen. (2) β, Triple; Mags. A 2, B 4, C 10; A–B, Pos. 25°, Dist. 13″; A–C, Pos. 89°, Dist. 0″.9. (3) ν (2° due east of β), Quadruple; Mags. A 4, B 5, C 7, D 8; A–B, Pos. 7°, Dist. 0″.8; A–C, Pos. 337°, Dist. 41″; C–D, Pos. 47°, Dist. 2″.4. A beautiful object. (4) ξ Scorpii (8½° due north from β), Triple; Mags. A 5, B 5.2, C 7.5; A–B, (Binary) Pos. 200°, Dist. 1″.4; ½(A + B) to C, Pos. 65°, Dist. 7″.3. μ^1 is a spectroscopic binary (Art. 465*).

Clusters. (1) M. 80, A.R. $16^h\ 10^m$; Dec. S. 22° 42′; half-way between α and β; one of the finest clusters known. (2) M. 4, A.R. $16^h\ 16^m$; Dec. S. 26° 14′; 1½° west of α; not so fine as the preceding.

Norma lies west of Scorpio, between it and *Lupus*, while *Ara* lies due south of η and θ. Both are small and of little importance, at least to observers in our latitudes.

48. Ophiūchus (*or Serpentarius*) and **Serpens** (Map III.).—Ophiuchus means the "serpent-holder." The giant is represented as standing with his feet on Scorpio, and grasping the "serpent," the head of which is just south of Corona Borealis, while the tail extends nearly to Aquila. The two constellations therefore are best treated together. The head of Serpens is marked by a group of small stars 20° due east of Arcturus, and 10° south of Corona. β and γ are the two brightest stars in the group, their magnitudes three and a half and four. δ lies 6° southwest of β, and there the serpent's body bends southeast through α and ϵ Serpentis (see map) to δ and ϵ Ophiuchi in the giant's hand. The line of these five stars carried upwards passes nearly through ϵ Boötis, and downwards

Digitized by Google

through ζ Ophiuchi. A line crossing this at right angles, nearly midway between ε Serpentis and δ Ophiuchi, passes through μ Serpentis on the southwest, and λ Ophiuchi to the northeast. The lozenge-shaped figure formed by the lines drawn from α Serpentis and ζ Ophiuchi to the two stars last mentioned forms one of the most characteristic configurations of the summer sky.

α Ophiuchi (2½ magnitude) is easily recognized in connection with α Herculis, since they stand rather isolated, about 6° apart, on the line drawn from Arcturus through the head of Serpens, and produced as far again. α Ophiuchi is the eastern and the brighter of the two. It forms with Vega and Altair a nearly equilateral triangle. β Ophiuchi lies about 9° southeast of α; and 5° east and a little south of β are five small stars in the Milky Way, forming a V with the point to the south, much like the Hyades of Taurus. They form the head of the now discredited constellation "Poniatowski's Bull" (*Taurus Poniatovii*), proposed in 1777.

49. *Names.* α Ophiuchi, *Ras Alaghue;* β, *Cebalrai;* δ, *Yed;* λ, *Marfic:* α Serpentis, *Unukalhai;* θ, *Alya.*

Double Stars. (1) λ Ophiuchi, Binary; period, 234 years; Mags. 4, 6; Pos. (1905) 60°; Dist. 1″.2. (2) 70 Ophiuchi (the middle star in the eastern leg of the V of Poniatowski's Bull), Binary; period, 93 years; Mags. 4.5, 7; Pos. (1905) 180°; Dist. 2″. The position angle changes very rapidly just now, and the star is too close to be resolved by a small instrument. (3) δ Serpentis, Mags. 4, 5; Pos. 185°; Dist. 3″.6; very pretty. (4) θ Serpentis, Mags. 4, 4.5; Pos. 104°; Dist. 21″. (5) ν Serpentis (4° N.E. of η Ophiuchi), Mags. 4.5, 9; Pos. 31°; Dist. 51″; sea-green and lilac.

Clusters. (1) M. 23, A.R. 17ʰ 50ᵐ; Dec. S. 19° 0′. Fine low-power field. (2) M. 12, A.R. 16ʰ 41ᵐ; Dec. S. 1° 45′. On the line between β and ε Ophiuchi, one-third of the way from ε. (3) M. 10, A.R. 16ʰ 51ᵐ; Dec. S. 3° 56′. On the line between β and ζ Ophiuchi, two-fifths of the way from ζ. (4) ℍ VIII. 72, A.R. 18ʰ 22ᵐ; Dec. N. 6° 29′. Pointed at by the eastern leg of the Poniatowski V. 8° from 70 Ophiuchi.

Digitized by Google

50. Hercules (Maps I. and III.). — This noble constellation lies next north of Ophiuchus, and is bounded on the west by Serpens, Corona, and Boötes, while to the east lie Aquila, Lyra, and Cygnus. On the north is Draco. The hero is represented as resting on one knee, with his foot on the head of Draco, while his head is close to that of Ophiuchus. The constellation contains no stars of the first or even of the second magnitude, but a number of the third. The most characteristic figure is the keystone-shaped quadrilateral formed by the stars ϵ, ζ, η, with π and ρ together at the northeast corner. It lies about midway on the line from Vega to Corona. The line $\pi\epsilon$, carried on 11°, brings us to β, the brightest star of the asterism; and γ and κ lie a few degrees farther along on the same line continued toward γ Serpentis. The angle $\epsilon\beta\alpha$ is a right angle opening towards Lyra. α is irregularly *variable*, besides being *double*.

51. *Names.* α, *Ras Algethi*; β, *Korneforos*.

Double Stars. (1) α, Mags. 3, 6; Pos. 119°; Dist. 4″.5; orange and blue. A very beautiful object for a 4-inch glass (Astr. Fig. 113). (2) ζ (the S.W. corner of the "Keystone"), Binary; period, 34 y. (Astr. Fig. 113); Mags. 3, 6.5; Pos. (1905) 190°; Dist. 1″.5. Rather difficult for a small instrument. (3) ρ ($2\frac{1}{2}$° east of π at the N.W. corner of the "Keystone"), Mags. 4, 5; Pos. 312°; Dist. 4″; white, emerald green. (4) δ (on the line from η through ϵ produced nearly its own length), Mags. 3, 8; Pos. 184°; Dist. 18″; white, light blue. Apparently an "optical pair"; the relative motion being rectilinear. (5) μ (nearly midway between Vega and α Herculis — see map), *Triple*; Mags. A 4, B 9.5, C 10; A, $\frac{B+C}{2}$, Pos. 246°; Dist. 31″. B–C, too faint and close for separation by any but large telescopes; Dist. about 1″; position angle rapidly changing — about 20° in 1890. (6) 95 Herculis (the N.W. corner of a little quadrilateral [sides 1° to 2°] of fourth and fifth mag. stars, on line from ρ through μ, produced two-thirds its length), Mags. 5.5 and 6; Pos. 262°; Dist. 6″; light green, cherry-red. Peculiar in showing contrast of color between *nearly equal components*.

Clusters. (1) M. 13, A.R. $16^h\ 37^m$; Dec. 36° 41′. Exactly on the western boundary of the Keystone, one-third the way from η towards

Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY

ζ. On the whole, the finest of all star clusters. (2) M. 92, A.R. $17^h\ 13^m$; Dec. 43° 16′ (6° north and a little west of ρ). Fine, but not equal to the other.

52. Lyra (Map IV.). — The great white or blue star VEGA sufficiently marks this constellation. It is attended on the east by two fourth-magnitude stars, ε and ζ, which form with it a little equilateral triangle having sides about 2° long. β and γ of the third magnitude (β is variable) lie about 8° southeast from Vega, $2\frac{1}{2}°$ apart.

Double Stars. (1) Vega itself has a small companion, 11th mag.; Pos. 160°; Dist. 48″. Only optically connected; the small star does not share the proper motion of the larger one, and has been used as a reference point in measuring Vega's parallax. (2) β, multiple; *i.e.*, it has three small stars near it, forming a very pretty object with a low power. (3) ϵ_1 and ϵ_2, Quadruple (the northern of the two which form the little triangle with α.) A sharp eye unaided by a telescope splits the star, and a small telescope divides both the components (see Astr. 468, and Fig. 113): ϵ_1 (or 4 Lyræ), Mags. 6, 7; Pos. 12°; Dist. 3″.2. ϵ_2 (or 5 Lyræ), Mags. 5.5, 6; Pos. 132°; Dist. 2″.5. $\epsilon_1\ \epsilon_2$, Pos. 173°; Dist. 207″. On the whole, the finest object of the kind. (4) ζ, Mags, 4, 6; Pos. 150°; Dist. 44″. (5) η (10′ E. of Vega), Mags. 4.5, 8; Pos. 85°; Dist. 28″; yellow, indigo. (6) δ; fine field for low powers.

Nebula. M. 57, the Annular Nebula. A.R. $18^h\ 49^m$; Dec. 32° 53′. Between β and γ, one-third of the way from β. (Art. 470.)

53. Cygnus (Maps I. and IV.). — This lies due east from Lyra, and is easily recognized by the cross that marks it. The bright star α ($1\frac{1}{2}$ magnitude) is at the top, and β (third magnitude) at the bottom, while γ is where the cross-bar from δ to ε intersects the main piece, which lies along the Milky Way from the northeast to the southwest. ζ is (nearly) on the prolongation of the line from γ through ε, not quite so far from ε as ε from γ.

Names. α, *Arided*, or *Deneb Cygni* (there are other *Denebs*; *e.g.*, *Deneb Kaitos* in Cetus); β, *Albireo*; γ, *Sadr*.

Digitized by Google

Double Stars. (1) β, Mags. 3.5, 7; Pos. 56°; Dist. 35″; orange, smalt blue. This is the finest of the colored pairs for a small telescope. (2) μ (as far beyond ζ as ζ is east of ϵ, at the tip of the eastern wing), Mags. 5 and 6; Pos. 118°; Dist. 3″.8. (3) χ (one-third of the way from β towards γ), Mags. 5 and 9; Pos. 73°; Dist. 26″; yellow and blue. (4) 61 Cygni (easily found by completing the parallelogram of which α, γ, and ϵ are the other three corners. σ and τ form a little triangle with 61, which is the faintest of the three), Mags. 5.5, 6; Pos. (1905) 127°; Dist. 22″. This is the star of which Bessel measured the parallax in 1838 (Astr. 521), — apparently our *second nearest* neighbor.

δ is also a fine double, but too difficult for an instrument of less than six inches' aperture.

Clusters. (1) M. 39, A.R. 21^h 28^m; Dec. 47° 54′ (about 3° north of ρ; ρ itself (fourth mag.) being found by drawing a line from δ through α, and carrying it an equal distance beyond. (2) ℍ VIII. 56, A.R. 20^h 19^m; Dec. 40° 20′. Beautiful group, ½° north and a little east of γ. The bright spots in the Milky Way all through Cygnus afford beautiful fields for a low power.

54. Vulpecula et Anser (Map IV.). — This little constellation is one of those originated by Hevelius, and has obtained more general recognition among astronomers than most of his creations. It lies just south of Cygnus, and is bounded to the south by Delphinus, Sagitta and Aquila.

It has no conspicuous stars, but contains one very interesting telescopic object, — the "Dumb-Bell Nebula," — M. 27, A.R. 19^h 54^m; Dec. 22° 23′. On a line from γ Lyræ through β Cygni, produced as far again, where this line intersects another drawn from α Aquilæ through γ Sagittæ, 3½° north and half a degree east of the latter star.

55. Sagitta (Map IV.). — This little asterism, though very inconspicuous, is one of the old 48. It lies south of Vulpecula, and the two stars α and β, which mark the feather of the arrow, lie nearly midway between β Cygni and Altair, while its point is marked by γ, 5° farther east and north.

Double Stars. (1) ζ ($\frac{3}{4}$° N.W. of δ, the middle star in the shaft of the arrow), Mags. 5.5, 9; Pos. 312°; Dist. 8″.6: the larger star is itself close double, distance about ½″, making an interesting triple system.

Digitized by Google

56. Aquila (Map IV.).—This constellation lies on the celestial equator, east of Ophiuchus and north of Sagittarius and Capricornus. It is bounded on the east by Aquarius and Delphinus, and on the north by Sagitta. Its characteristic configuration is that formed by ALTAIR (the standard first-magnitude star), with γ to the north and β to the south. It lies about 20° south of β Cygni, and forms a fine triangle with Vega and α Ophiuchi.

Double Star. (1) π Aquilæ (1½° N.E. of γ), Mags. 6 and 7; Pos 120°; Dist. 1″.5. Good test for 3½-inch glass.

Cluster. M. 11, A.R. 18ʰ 45ᵐ; Dec. S. 6° 24′. A fine fan-shaped group of stars in the Milky Way. A line carried from Altair through δ Aquilæ (see map), and prolonged once and a half as far again, will find it about 4° S.W. of λ.

The southern part of the region allotted to Aquila on our maps has been assigned to *Antinoüs*. This constellation was recognized by some even in Ptolemy's time; but he declined to adopt it. Hevelius appropriated the eastern portion of "Antinoüs" for his constellation of "*Scutum Sobieski*," and M. 11 falls just within its limits.

57. Sagittarius (Map IV.).—This, the tenth of the zodiacal constellations, is bounded north by Aquila and Ophiuchus, west by Scorpio and Ophiuchus (though Bode and some other authorities crowd in a piece of "Telescopium" between it and Scorpio), south by Corona Australis, Telescopium, and Indus, and east by Microscopium and Capricornus. It contains no stars of the first magnitude, but a number of the 2½ and third.

The most characteristic configuration is the little inverted "milk dipper" formed by the five stars, λ, ϕ, σ, τ, and ζ, of which the last four form the bowl, while λ (in the Milky Way) is the handle. δ, γ, and ϵ, which form a triangle right-angled at δ, lie south and a little west of λ, the whole eight together forming a very striking group. There is a curious disregard of any apparent principle in the lettering of the stars of this constellation; α and β are stars not exceeding in brightness

Digitized by Google

the fourth magnitude, about 4° apart on a north and south line and lying some 15° south and 5° east of ζ (see map). The Milky Way in Sagittarius is very bright, and complicated in structure, full of knots and streamers, and dark pockets.

Names. λ, *Kaus Borealis*; δ, *Kaus Media*; ϵ, *Kaus Australis*; σ, *Sádira*. This star is strongly suspected of irregular variability.

Double Stars. (1) μ^1 (7° N.W. of λ; on the line from ζ through ϕ produced), Triple; Mags. A 3.5, B 9.5, C 10; A–B, Pos. 315°, Dist. 40″; A–C, Pos. 114°, Dist. 45″.

Clusters and Nebulæ. (1) M. 22, A.R. $18^h\ 29^m$; Dec. S. 24° 0′ (3° N.W. of λ, and midway between μ and σ). Capital object for a 4-inch telescope. (2) M. 25, A.R. $18^h\ 25^m$; Dec. S. 19° 10′ (7° north and 1° east of λ; visible to naked eye). (3) M. 8; A.R. $17^h\ 57^m$; Dec. S. 24° 21′ (a little south of the line $\phi\lambda$ produced, and as far from λ as λ from ϕ; also visible to naked eye). (4) ♅ IV. 41, The *Trifid Nebula*, A.R. $17^h\ 55^m$; Dec. S. 23° 2′ (1¼° north of M. 8, and almost exactly on the line $\phi\lambda$ produced). A very beautiful and interesting object.

58. Capricornus (Map IV.). — This, the eleventh of the zodiacal constellations, follows Sagittarius on the east. It has Aquarius and Aquila (Antinoüs) on the north, Microscopium and Piscis Austrinus on the south, and Aquarius on the east. It has no bright stars, but the configuration formed by the two α's (α_1 and α_2) with each other and with β, 3° south, is characteristic and not easily mistaken for anything else. The two α's, a pretty "double" to the naked eye, lie on the line from β Cygni (at the foot of the cross) through Altair, produced about 25°. On the line $\alpha\beta$, about 3° distant, lies ρ (of the fourth magnitude), with two other small stars near it. From this a line 20° long, carried due east through θ and ι (of the fourth magnitude), brings the eye to γ and δ of the third, the latter marking the constellation's eastern limit.

Names. α, *Algiedi* (*prima* and *secunda*); δ, *Deneb Algiedi*.

Double Stars. (1) α_1 and α_2 (pretty with a very low power), Mags. 3 and 4; Dist. 6′ 13″. α_2 has also a very faint companion, invisible

Digitized by Google

with any telescope of less than 6-inch aperture; Pos. 150°; Dist. 7″.5. The companion is itself double; Dist. about 1″; Pos. 240°. (2) β, Mags. 3.5, 7; Pos. 267°; Dist. 3′ 25″. The companion is also a close and difficult double. (3) ρ (the northern star in the little triangle it forms with π and o), Mags. 5, 9; Pos. 177°; Dist. 3″.8. (4) π (the S.W. one in the same triangle), Mags. 5, 9; Pos. 146°; Dist. 3″.5.

Nebula. M. 30, A.R. $21^h\ 34^m$; Dec. S. 23° 42′ (about 1° west and a little north of 41 Capricorni, a fifth-magnitude star, 7° south of γ Capricorni).

59. Delphinus (Map IV.). — This little asterism is ancient, and unmistakably characterized by the rhombus of third-magnitude stars known as "Job's Coffin." It lies about 15° northeast of Altair, bounded north by Vulpecula and west by Aquila. There are a few stars visible to the naked eye in addition to the four that form the rhombus. *Epsilon*, about 3° southwest, is the only conspicuous one.

Names. *α, Svalocin; β, Rotanev.* These were given in joke by Nicolaus Cacciatore, a Sicilian astronomer, about 1800. The letters of the two names *reversed* make Nicolavs Venator; Venator being the translation of the Italian "Cacciatore," which means "*Hunter.*" The joke is good enough to keep.

Double Stars. (1) γ (at the N.E. angle of the rhombus), Mags. 4, 7; Pos. 271°; Dist. 11″.3. (2) β, a very close and rapid binary, beyond the reach of all but large telescopes. It has, however, two little companions, distant about 30″.

60. Equuleus (Map IV.). — This little constellation is still smaller than the Dolphin, and contains no such characteristic star group. It lies about 20° due east of Altair, and 10° S.E. of Delphinus (see map).

Double Stars. (1) ϵ, Mags. 5, 7.5; Pos. 73°; Dist. 11″. The larger star is also close double; Mags. 5.5. and 7; Pos. 290°; Dist. 0″.9. Perhaps resolvable with a 4-inch telescope.

61. Lacerta (Maps I. and IV.). — This is one of Hevelius's modern constellations, lying between Cygnus and Andromeda, with no stars above the 4½ magnitude. It contains a few telescopic objects, but nothing suited to our purpose.

Digitized by Google

62. Pēgăsus (not Pegas'us) (Map IV.). — This covers an immense space which is bounded on the north by Andromeda and Lacerta, on the west by Cygnus, Vulpecula, Delphinus, and Equuleus, on the south by Aquarius and Pisces, and on the east by Pisces and Andromeda. Its most notable configuration is "the great square," formed by the second-magnitude stars α, β, and γ Pegasi, in connection with α Andromedæ (sometimes lettered δ Pegasi) at its northeast corner. The stars of the square lie in the body of the horse, which has no hindquarters. The line drawn from α Andromedæ through α Pegasi, and produced about an equal distance, passes through ξ and ζ in the animal's neck, and reaches θ (third magnitude) in his ear. *Epsilon*, 8° northwest of θ, marks his nose. The forelegs are in the northwestern part of the constellation just east of Cygnus, and are marked, one of them by the stars η and π, the other by ι and κ.

Names. α, *Markab*; β, *Scheat*; γ, *Algenib*; ϵ, *Enif*.

Double Star. κ, Mags. 4, 11; Pos. 302°; Dist. 12″. The large star is also itself an extremely close double; Dist. 0″.3; (pointed at by the northern edge of the "square," at a distance one and a quarter times its length.)

Cluster. M. 15, A.R. $21^h\ 24^m$; Dec. 11° 38′ (on the line from θ through ϵ, produced half its length, and just west of a sixth-magnitude star).

63. Aquarius (Map IV.). — This, the twelfth and last of the zodiacal constellations, extends more than $3\frac{1}{2}$ hours in right ascension, covering a considerable region which by rights ought to belong to Capricornus. It is bounded north by Delphinus, Equuleus, Pegasus, and Pisces; west by Aquila and Capricornus; south by Capricornus and Piscis Austrīnus, and east by Cetus. The most notable configuration is the little Y of third and fourth magnitude stars which marks the "water jar" from which Aquarius pours the stream that meanders down to the southeast and south for 30°, till it

Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY

eaches the Southern Fish. The middle of the Y is about 18° outh and west of α Pegasi, and lies almost exactly on the elestial equator. A line drawn west and a little south from ν (the westernmost star of the Y) to α Capricorni, passes through β (third magnitude) at one-third of the way, and through μ and ε (fourth and 3½) two-thirds of the way. α (third magnitude) lies 4° west and a little north of γ. δ (third magnitude) lies about half-way between the Y and Fomalhaut in the Southern Fish, 3° or 4° east of the line that joins them.

Names. α, *Saad el Melik*; β, *Saad el Suna*, δ, *Skat.*

Double Stars. (1) ζ (the central star of the Y), Mags. 4, 4.5; Pos. 332°; Dist. 3″.6; pretty and easy. (2) 12 Aquarii (7° due west of β, and the brightest star in the vicinity), Mags. 5.5, 8.5; Pos. 190°; Dist. 2″.8; yellowish white and light blue.

Clusters and Nebulæ. (1) M. 2; A.R. 21ʰ 17ᵐ; Dec. S. 1° 22′ (on the line drawn from ζ through α, produced one and a quarter times its own length). (2) H IV. 1, A.R. 20ʰ 58ᵐ; Dec. S. 11° 50′ (nearly on the line from α through β, produced its own length, and 1⅓° west of ν; fifth magnitude); planetary nebula, bright and vividly *green.*

64. Piscis Austrīnus (*or Australis*) (Map IV.).—This small constellation, lying south of Aquarius and Capricornus, presents little of interest. It has one bright star, FOMALHAUT (pronounced *Fomalhawt*), of the 1½ magnitude, which is easily recognized from its being nearly on the same hour-circle with the western edge of the great square of Pegasus, 45° to the south of α, and solitary, having no star exceeding the fourth magnitude within 15° or 20°. It contains no telescopic objects available for our purpose.

South of it, barely rising above the southern horizon, lie the constellations of *Microscopium* and *Grus.* The former is of no account. The latter is a conspicuous constellation in the southern hemisphere, and its two brightest stars, α and β, of the second magnitude, rise high enough to be seen in latitudes south of Washington. They lie about 20° south and west of Fomalhaut.

Digitized by Google

LIST OF CONSTELLATIONS, SHOWING THEIR POSITION IN THE HEAVENS, AND THE NUMBER OF STARS IN EACH.

(The zodiacal constellations are denoted by italics, non-Ptolemaic constellations by an asterisk.)

R. A. \ Decl.	+ 90° to + 50°.	+ 50° to + 25°.	+ 25° to 0°.	0° to − 25°.	− 25° to − 50°.	− 50° to − 90°.
h. h.						
I, II,	Cassiopeia, 46	Andromeda, 18 Triangulum, 5	*Pisces*, 18 *Aries*, 17	Cetus, 32	Phœnix, 32 *App. Sculp. 13	Phœnix, *bis.* Hydrus, 18
III, IV,	– –	Perseus, 40	*Taurus*, 58	Eridanus, 64	(Eridanus, *bis.*)	*Horologium, 11 *Reticulum, 9
V, VI,	*Camelopardalis, 36	Auriga, 35	Orion, 37 *Gemini*, 28	Lepus, 18	*Columba, 15	*Dorado, 16 *Pictor, 14 *Mons Mensæ, 12
VII, VIII,	– –	*Lynx, 28	Canis Minor, 6 *Cancer*, 15	Canis Major, 27 *Monoceros, 12	Argo-Navis, 133	Argo-Navis, *bis.* (Puppis) *Piscis Volans, 9
IX, X,	– –	*Leo Minor, 15	*Leo*, 47	Hydra, 49 *Sextans, 3	– –	Argo-Navis (Vela)
XI, XII,	Ursa Major, 53	– –	*Coma Ber. 20	Crater, 9 Corvus, 8	Centaurus, 54	Argo-Navis (Carina) *Chameleon, 13
XIII, XIV,	– –	*Canes Venat. 15 Boötes, 35	– –	*Virgo*, 39	Lupus, 34	Centaurus, *bis.* *Crux, 13 *Musca, 15
XV, XVI,	Ursa Minor, 23	Corona Bor. 19 Hercules, 65	Serpens, 23	*Libra*, 23	Norma,* 14	*Circinus, 10
XVII, XVIII,	Draco, 80	Lyra, 18	Aquila, 37 Sagitta, 5	*Scorpio*, 34 Ophiuchus, 46	Ara,* 15	*Triangul. Aust. 11 *Apus, 15
XIX, XX,	– –	Cygnus, 67	*Vulpecula, 23 Delphinus, 10	*Sagittarius*, 38	Corona Austr. 7	*Telescopium, 16 Pavo, 37 *Octans, 22
XXI, XXII,	Cepheus, 44	*Lacerta, 13	Equuleus, 5	*Capricornus*, 22	Piscis Austr. 16	*Indus, 15 *Octans,
XXIII, XXIV,	– –	– –	Pegasus, 43	*Aquarius*, 25	*Grus, 30	*Toucana 22 *Octans.

Digitized by Google


Digitized by Google

Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY

9 P.M. MAP I.



Digitized by Google

Digitized by Google

Digitized by Google



Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY



Digitized by Google

Digitized by Google



Digitized by Google



Digitized by Google



Digitized by Google

Digitized by Google

Digitized by Google



Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY
BERKELEY

MAP IV.



Digitized by Google

Digitized by Google

G
P.
UNIVERSITY OF CALIFORNIA LIBRARY
BERKELEY

Digitized by Google

Digitized by Google

UNIVERSITY OF CALIFORNIA LIBRARY
BERKELEY

Digitized by Google

Digitized by Google

Digitized by Google

Digitized by Google

Digitized by Google